

[Journal of Physical Organic Chemistry](#)
Volume 13, Issue 8 , Pages 443 – 451

Research Article

Regiochemistry and mechanism of oxidation of *N*-benzyl-*N*-alkylhydroxylamines to nitrones

Azfar Hassan, Mohammed I. M. Wazeer, Mohammed T. Saeed, Mohammad N. Siddiqui, Sk. Asrof Ali *

Chemistry Department, King Fahd University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia

email: Sk. Asrof Ali (shaikh@kfupm.edu.sa)

*Correspondence to Sk. Asrof Ali, Chemistry Department, King Fahd University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia

Keywords

hydroxylamines; oxidation; mercury(II) oxide; *p*-benzoquinone; isotope effect; nitrones; isomerization; kinetics

Abstract

The oxidation of various *N*-(*o*-, *m*-, *p*-substituted benzyl)-*N*-alkylhydroxylamines and their dideuteriobenzyl (PhCD₂) counterparts was carried out using mercury(II) oxide and *p*-benzoquinone (*p*-BQ) as oxidants. An overwhelming preference for the formation of conjugated nitrones is observed in the oxidation of *N*-benzyl-*N*-isopropylhydroxylamines. Considerable intra- and intermolecular kinetic isotope effects and negative ρ values in the Hammett plots point towards a mechanistic pathway that involves electron transfer from nitrogen to the oxidant followed by hydrogen abstraction. The conformation of unstable (*E*)-nitrones, which readily isomerize to the more stable (*Z*)-nitrones, is deduced from ¹H NMR data. The *E* ⇌ *Z* isomerization was found to be a bimolecular process. Copyright © 2000 John Wiley & Sons, Ltd.

Received: 21 September 99; Revised: 15 December 99; Accepted: 9 February